

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

DORLAND'S ILLUSTRATED

Medical Dictionary

Twenty-sixth Edition

W. B. SAUNDERS COMPANY Philadelphia London Toronto
Mexico City Sydney Tokyo

**The Library of Congress Cataloged the First Issue
of this Serial as follows:**

Dorland's illustrated medical dictionary. [1st] — ed.

Philadelphia, Saunders, 1900—

illus. (part col.) 23–25 cm.

Title varies: 1st–22d ed., The American illustrated medical dictionary.

- | | |
|---------------------------|-------------------------------------|
| I. Medicine—Dictionaries. | I. Dorland, William Alexander |
| Newman, 1864–1956. | II. Title: The American illustrated |
| medical dictionary. | |

R121.D73 610.3 0–6383 rev 4*

Library of Congress [r65i²7] MARC–S

© 1981 by W. B. Saunders Company

Copyright 1900, 1901, and 1903 by W. B. Saunders and Company. Copyright 1906, 1909, 1911, 1913, 1915, 1917, 1919, 1921, 1923, 1925, 1927, 1929, 1932, 1935, 1938, 1941, 1944, 1947, 1951, 1957, 1965, and 1974 by W. B. Saunders Company.

Copyright under the Uniform Copyright Convention. Simultaneously published in Canada. All Copyright Renewals Registered.

Derechos reservados conforme a la ley para la Republica Mexicana.

All Rights Reserved. This book is protected by copyright. No part of it may be duplicated or reproduced in any manner without written permission from the publisher. Made in the United States of America. Press of W. B. Saunders Company.

Some of the words appearing in the Dictionary are proprietary names (trademarks) even though no reference to this fact is made in the text. The appearance of any name without designation as a trademark is therefore not to be regarded as a representation by the editors or publisher that it is not a trademark or is not the subject of proprietary rights.

The use of portions of the text of the *United States Pharmacopeia*, Twentieth Revision, official from July 1, 1980, of the *National Formulary*, Fifteenth Edition, official from July 1, 1980, and of *USAN and the USP Dictionary of Drug Names 1981* is by permission received from the Board of Trustees of the United States Pharmacopeial Convention, Inc. The said Convention is not responsible for any inaccuracy of quotation, or for any false or misleading implication that may arise by reason of the separation of excerpts from the original context or by obsolescence resulting from publication of a supplement.

Listed here are the latest translated editions of this book together with the languages for the translations and the publishers.

Japanese (25th Edition)—Hirokawa Publishing Company, Tokyo, Japan

Spanish (25th Edition) (Adaptation)—El Ateneo, Buenos Aires, Argentina

Braille edition (24th Edition)—American Printing House for the Blind, Louisville, Kentucky

ISBN 0-7216-3150-9 Standard
ISBN 0-7216-3151-7 Indexed

Library of Congress Catalog Card Number: 78-50050

Last digit is the print number: 9 8 7 6 5 4 3 2

yellowish white crystals or powder, usually used in combination with other sulfonamides. See *trisulfapyrimidines oral suspension*, under *suspension*. Called also *sulfamethyldiazine*.

sulfameter (sul'fah-me'ter) chemical name: 4-amino-*N*-(5-methoxy-2-pyrimidinyl)benzenesulfonamide. A long-acting sulfonamide, $C_{11}H_{11}N_3O_3S$, occurring as a fine, white to yellowish-white, powder; used as an antibacterial, especially in the treatment of acute and chronic urinary tract infections, administered orally.

sulfamethazine (sul'fah-meth'ah-zën) [USP] chemical name: 4-amino-*N*-(4,6-dimethyl-2-pyrimidinyl)benzenesulfonamide. A sulfonamide, $C_{12}H_{14}N_2O_3S$, occurring as a white to yellowish white powder; used as an antibacterial in a variety of infections, usually, in the United States, in combination with other sulfonamides. It is administered orally. Called also *sulfadimidine*.

sulfamethizole (sul'fah-meth'i-zöl) [USP] chemical name: *N*-(5-methyl-1,3,4-thiadiazol-2-yl)sulfanilamide. A compound, $C_9H_{10}N_4O_2S_2$, occurring as white crystals or powder, used as an antibacterial agent mainly in the treatment of infections of the urinary tract. Called also *sulfamethylthiadiazole*.

sulfamethoxazole (sul'fah-meth-oks'ah-zöl) [USP] chemical name: 4-amino-*N*-(5-methyl-3-isoxazolyl)benzenesulfonamide. A sulfonamide, $C_{10}H_{11}N_3O_3S$, occurring as a white to off-white, crystalline powder; used as an antibacterial, especially for the prophylaxis and treatment of acute urinary tract infections and of pyoderma and infections of wounds and soft tissues, administered orally.

sulfamethoxypyridazine (sul'fah-meth-ok'se-pi-rid'ah-zën) chemical name: *N*-(6-methoxy-3-pyridazinyl)sulfanilamide. A compound, $C_{11}H_{12}N_4O_3S$, occurring as a white or yellowish white, crystalline powder, used as an antibacterial agent in the treatment of infections of the urinary tract and other infections.

sulfamethyldiazine (sul'fah-meth'il-di'ah-zën) sulfamazine.

sulfamethylthiadiazole (sul'fah-meth'il-thi'ah-di'ah-zöl) sulfamethizole.

Sulfamezathine (sul'fah-mez'ah-thën) trademark for a preparation of sulfamethazine.

sulfamido (sul-fam'i-do) one of a group of compounds containing an aminosulfone group $SO_2 \cdot NH_2$.

sulfamine (sul-fam'in) the univalent radical, $-SO_2NH_2$.

sulfamonomethoxine (sul'fah-mon'o-më-thoks'ën) chemical name: *N*-(6-methoxy-4-pyrimidinyl)sulfanilamide; an antibacterial, $C_{11}H_{11}N_4O_3S$.

sulfamoxole (sul'fah-moks'öl) chemical name: *N*-(4,5-dimethyl-2-oxazolyl)sulfanilamide; an antibacterial, $C_{11}H_{13}N_3O_3S$.

Sulfamylon (sul'fah-mi'lon) trademark for preparations of mafenide.

sulfanilamide (sul'fah-ni'l'ah-mid) chemical name: *p*-amino-benzenesulfonamide. A potent antibacterial compound, $NH_2 \cdot C_6H_4 \cdot SO_2NH_2$, the first of the sulfonamides discovered. Formerly used in the treatment of various infections, it has been replaced by more effective and less toxic derivatives, and by antibiotics. Called also *prontosil album*.

sulfanilate (sul-fan'i-lät) a salt of sulfanilic acid.

sulfanitrane (sul'fah-ni'tran) chemical name: 4'-(*p*-nitrophenyl)sulfamoylacetanilide. A substance, $C_{14}H_{13}N_3O_5S$, used as an antibacterial and as a coccidiostatic agent in poultry.

sulfanuria (sul'fah-nu're-ah) anuria resulting from the use of sulfonamide drugs.

sulfapyridine (sul'fah-pir'i-dën) [USP] chemical name: *N*-(2-pyridyl)sulfanilamide. An antibacterial compound, $C_{11}H_{11}N_3O_2S$, occurring as white or faintly yellowish white granules, crystals, or powder; used as an oral suppressant for dermatitis herpetiformis. It was formerly used in the treatment of pneumonia and streptococcal infections.

sulfaquinoxaline (sul'fah-kwin-ok'sah-lën) chemical name: *N*-(2-quinoxalyl)sulfanilamide. An antibacterial, $C_{14}H_{12}N_2O_5S$, used in veterinary medicine as a coccidiostat, and in the treatment of fowl cholera, fowl typhoid, infectious enteritis of swine, shipping dysentery of lambs, and foot rot of cattle.

sulfarsphenamine (sul'ar-sfen'ah-min) the disodium salt of dihydroxy-diaminoarsenobenzenemonomethylene sulfonate, $NH_2(OH)C_6H_3 \cdot As \cdot AsC_6H_3(OH)NH_2 \cdot CH_2 \cdot SO_2 \cdot ONa$. It contains 18-20 per cent of arsenic and was formerly used in the treatment of syphilis. It differs from neoarsphenamine in having two side chains instead of one and in that the sulfur has a valence of four instead of two.

sulfasalazine (sul'fah-sal'ah-zën) [USP] chemical name: 2-hydroxy-5-[[4-[(2-pyridinylamino)sulfonyl]phenyl]azo]benzoic acid. An antibacterial sulfonamide derivative, $C_{18}H_{14}N_4O_5S$, occurring as a bright yellow to brownish yellow powder; used orally in the treatment of mild to moderate ulcerative colitis and as adjunctive therapy in severe ulcerative colitis due to susceptible organisms, administered orally. Called also *salazosulfapyridine* and *salicylazosulfapyridine*.

Sulfasuxidine (sul'fah-suk'si-dën) trademark for preparations of succinylsulfathiazole.

sulfatase (sul'fah-täs) an enzyme that catalyzes the hydrolysis of various sulfuric acid esters into sulfuric acid and alcohol.

sulfate (sul'fät) [L. *sulphas*] any salt of sulfuric acid. **acid s.**, one in which only one half of the hydrogen of the sulfuric acid is replaced; a bisulfate. **basic s.**, one in which the normal sulfate of the base is combined with a hydroxide of the same base; a subsulfate. **chondroitin s.**, see *chondroitin*. **conjugated s's**, aromatic substances, such as phenol, scatoxyl, and indoxyl, which occur in the urine along with mineral sulfates. **cupric s.** [USP], the pentahydrate sulfate salt of copper, $CuSO_4 \cdot 5H_2O$, occurring as deep blue, triclinic crystals or as blue crystalline granules or powder, which is a powerful emetic; used orally as an antidote to phosphorus poisoning. Topical application of a 1 per cent solution is used in the treatment of phosphorus burns of the skin. It is also used as a catalyst with iron in the treatment of iron deficiency anemia. In 1:1,000,000 concentration it is used to prevent growth of algae in ponds, reservoirs, and swimming pools. Called also *blue vitriol*, *copper sulfate*, and *bluestone*. **dermatan s.**, chondroitin sulfate B; see *chondroitin*. **etheral s's**, conjugated s's. **ferrous s.** [USP], pale bluish green odorless crystals or granules, $FeSO_4 \cdot 7H_2O$, used orally in treatment of iron deficiency anemia. Called also *copperas*, *green vitriol*, *iron protosulfate*, and *iron sulfate*. **ferrous s. dried** [USP], a grayish white powder, $FeSO_4 \cdot H_2O$, used as a hematinic. **mineral s's**, sulfates in the urine which are combinations of sulfuric acid with mineral substances such as sodium, potassium, calcium, and magnesium. **neutral s.**, **normal s.**, one in which all the hydrogen of the sulfuric acid is replaced. **preformed s's**, mineral s's.

sulfatemia (sul'fät-e'me-ah) the presence of sulfates in the blood.

Sulfathalidine (sul'fah-thal'i-dën) trademark for phthalylsulfathiazole.

sulfathiazole (sul'fah-thi'ah-zöl) chemical name: *N*-(2-thiazolyl)sulfanilamide. A compound, $C_9H_9N_3O_2S_2$, once widely used as an antibacterial agent but replaced by less toxic sulfonamides and antibiotics. Called also *M & B 760*, *norsulfazole*.

sulfatide (sul'fah-tid) one of a class of cerebroside sulfuric esters; they are found largely in the medullated nerve fibers, and may accumulate in the white matter of the brain in metachromatic leukodystrophy.

sulfazamet (sul-fah'zah-met) chemical name: *N*-(3-methyl-1-phenylpyrazol-5-yl)sulfanilamide; an antibacterial, $C_{14}H_{15}N_3O_2S$.

sulfhemoglobin (sul'he-mo-glo'bin) sulfmethemoglobin.

sulfhemoglobinemia (sul'he-mo-glo'bin-e'me-ah) the presence of sulfmethemoglobin in the blood.

sulphydrate (sul-fi'drät) any compound of a base with sulphydric acid or, more correctly, with the radical sulphydryl, SH, or hydrogen sulfide.

sulphydryl (sul-fi'dril) the univalent radical, $-SH$.

sulfide (sul'fid) any binary compound of sulfur; a compound of sulfur with another element or radical or base. **mercuric s.**, a brilliant scarlet powder, HgS, formerly used in the treatment of syphilis.

sulfindigotate (sul-fin'di-go-tät) any salt of sulfindigotic acid.

sulfipyrazone (sul'fin-pi'rah-zön) [USP] chemical name: 1,2-diphenyl-4-[2-(phenylsulfinyl)ethyl]-3,5-pyrazolidinedione. A sulfoxide analogue of phenylbutazone, $C_{22}H_{19}N_2O_3S$, used as a uricosuric agent in treatment of gout.

sulfinyl (sul'fi-nil) the bivalent radical, $-SO-$.

sulfisomidine (sul-fi-som'i-dën) chemical name: *N*-(2,6-dimethyl-4-pyrimidinyl)sulfanilamide. A compound, $C_{12}H_{14}N_4O_2S$, closely related to sulfamethazine, occurring as a white or creamy-white powder, used as an antibacterial agent in the treatment of systemic and urinary tract infections. Called also *sulfadimetine*.

sulfisoxazole (sul'fi-sok'sah-zöl) [USP] chemical name: 4-amino-*N*-(3,4-dimethyl-5-isoxazolyl)benzenesulfonamide. A short-acting sulfonamide, $C_{11}H_{13}N_3O_3S$, occurring as a white to slightly yellowish, crystalline powder; used as an antibacterial in the treatment of a wide variety of infections, administered orally. Called also *sulfafurazole*. **acetyl s.** [USP], a tasteless derivative of sulfisoxazole, having the same actions as the base. **s. diethanolamine**, **s. diolamine** [USP], a soluble salt of sulfisoxazole, administered parenterally.

sulfite (sul'fit) [L. *sulfis*] any salt of sulfurous acid. **s. oxidase**, an oxidoreductase that catalyzes the oxidation of sulfite (with O_2) to sulfate with release of H_2O_2 ; it is a hemoprotein containing molybdenum, occurring in the intermembrane space of mitochondria.

sulfmethemoglobin (sul'fet-he'mo-glo'bin) a greenish substance formed by treating blood with hydrogen sulfide or by the absorption of this gas from the intestinal tract; it is the cause of the greenish color seen in the abdominal walls and along the vessels of cadavers. Called also *sulfhemoglobin*.

sulfo- a prefix used in naming chemical compounds, indicating the presence of divalent sulfur or of the group SO_2OH .

sulfoacid (sul'fo-as'id) sulfonic acid.

BIOCHEMICALS
ORGANIC COMPOUNDS
DIAGNOSTIC REAGENTS

TOLL FREE ORDERING

**Ordering
Hours**

Monday through Friday - 24 Hours a day
Saturday, Sunday and Holidays -
6:00 a.m. to 5:00 p.m. (CST)

Telephone

Call Collect.....314-771-5750

Bulk Inquiries Call SAF
See Page 15 for Details

FAX

For TWX, TELEX, Cable or Collect FAX Calls see page 3

CUSTOMER/TECHNICAL SERVICE

In Japan

Please callTokyo (03) 3258 0155

Should you have any problems getting excellent service at the above numbers, please call COLLECT (anytime Day or Night, including Weekends and Holidays) 314-531-9500.

SIGMA

ALPHABETICAL
LIST

BIOACTIVE
PEPTIDES

IMMUNO-
CHEMICALS

MOLECULAR
BIOLOGY

TISSUE
CULTURE

OTHER
PRODUCT
GROUPS

EQUIPMENT
BOOKS AND
SUPPLIES

DIAGNOSTIC
KITS AND
REAGENTS

SULFATASE(Aryl-sulfatase; Phenolsulfatase; Aryl-sulfate
sulfohydrolase; EC 3.1.6.1)**Sulfatase Unit Definition:** One unit will hydrolyze
1.0 μ mole of p-nitrocatechol sulfate per hr at
pH 5.0 at 37°C (30 min assay), unless otherwise
indicated below. **β -Glucuronidase Unit Definition:** One Sigma or
modified "Fishman" unit will liberate 1.0 μ g of
phenolphthalein from phenolphthalein glucuronic
acid per hr at pH 3.8 at 37°C (30 min assay),
unless indicated below.

[9016-17-5]

S 8504 (O-C)	Type IV: From Limpets	50 units	7.75
	(Patella vulgata)	100 units	12.35
	Lyophilized, essentially salt-free powder	500 units	42.25
		1,000 units	74.75
	Activity: 10-25 units per mg solid.	5,000 units	293.30

β -Glucuronidase Activity: 1,000-2,000 Sigma units per mg solid.

S 8629 (O-C)	Type V: From Limpets	100 units	14.50
	(Patella vulgata)	250 units	30.50
	Lyophilized, essentially salt-free powder	500 units	49.80
		1,000 units	89.05
	Activity: 5-15 units per mg solid.	5,000 units	348.60

β -Glucuronidase Activity: <2 Sigma units per mg solid.

S 1629 (O-C)	Type VI: From Aerobacter aerogenes	10 units	23.30
		50 units	102.10

Partially purified
50% glycerol solution (10-20 units per ml)
containing 0.01 M Tris, pH 7.5.
Activity: 2-5 units per mg protein (Biuret).
No detectable β -glucuronidase activity at pH 7.
Unit Definition: One unit will hydrolyze 1.0 μ mole
of p-nitrophenyl sulfate per min at pH 7.1 at 37°C.

S 9629 (O-C)	Type VII: From Abalone	100 units	11.40
	Entrails	500 units	40.40
	Lyophilized powder	1,000 units	68.05
		5,000 units	298.55
	Activity: 30-80 units per mg solid.		

β -Glucuronidase Activity: 400-800 Sigma units per mg solid.

S 9754 (O-C)	Type VIII: From Abalone	100 units	15.25
	Entrails	500 units	55.15
	Lyophilized powder	1,000 units	95.10
		5,000 units	412.30
	Activity: 20-40 units per mg solid.		

β -Glucuronidase Activity: <3.0 Sigma units per mg solid.

S 9626 (O-C)	Type H-1: From Helix pomatia (pfs)	5,000 units	10.80
		10,000 units	16.55
	Partially purified powder	100,000 units	90.35

Activity: 15-40 units per mg solid.
 β -Glucuronidase Activity: Approx. 300 units per mg solid at pH 5.0.

S 9751 (O-C)	Type H-2: From Helix pomatia (pfs)	1 ml	13.95
	Crude solution	2 ml	23.30
	Activity: 2,000-5,000 units per ml of solution.	5 ml	51.65
		10 ml	92.25
	β -Glucuronidase Activity: Approx. 100,000 Sigma units per ml at pH 5.0.	25 ml	202.35

(Continued)

(Continuation of)
SULFATASE

S 3009 (O-C)	Type H-5: From Helix pomatia	5,000 units	14.55
		10,000 units	24.80
	Lyophilized powder	100,000 units	173.95

A further purification of
S 9751 by gel filtration chromatography.
Sulfatase Activity: 15-40 units per mg solid.
 β -Glucuronidase Activity: 400-600 units per mg solid.

See also: Limpet Acetone Powder Page 625

SULFATHIAZOLE

(4-Amino-N-2-thiazolylbenzenesulfonamide)

S 9876 (RT)	Free Acid (pfs)	100 g	11.15
		250 g	21.35
	Crystalline	500 g	34.55
		1 kg	56.45
			FOB Sigma

[72-14-0] C₉H₉N₃O₂S₂ FW 255.3

S 0127 (RT)	Sodium Salt (pfs)	100 g	12.40
		250 g	23.30
	Crystalline	500 g	38.10
		1 kg	63.25
			FOB Sigma

[144-74-1] C₉H₈N₃O₂S₂Na
FW 277.3

SULFATIDES

S 1006 (O-C)	(Cerebroside sulfate)	5 mg	26.55
		10 mg	43.40
	From Bovine Brain	25 mg	86.05
		50 mg	154.25
	Also available as part of a kit.	100 mg	277.50

See: Standards and Controls
Section Page 1912
See also: Tissue Culture Media and Reagents
Page 1513

SULFHYDRYL CELLULOSESee under: Affinity Chromatography Media
Page 1731

S 9509 (RT)	(\pm)-SULFINPYRAZONE (pfs)	1 g	7.75
		5 g	19.40
		10 g	29.30
		25 g	51.90

[57-96-5] C₂₃H₂₀N₂O₃S FW 404.5

S 8259 (RT)	2,2'-SULFINYL-bis(4,6-DICHLORO-PHENOL) (pfs)	25 g	9.75
		100 g	25.60
	(Bithionol sulfoxide)	250 g	50.00
		1 kg	136.40

[844-26-8] C₁₂H₆O₃Cl₂S FW 372

S 7505 (RT)	SULFISOMIDINE (pfs)	25 g	9.45
		100 g	29.35
	(4-Amino-N-[2,6-dimethyl-4-pyrimidinyl]benzenesulfonamide)		
	Crystalline		

[515-64-0] C₁₂H₁₄N₄O₂S FW 278.3

S 6377 (O-S-C)	SULFISOXAZOLE (pfs)	25 g	5.15
		100 g	11.55
	(4-Amino-N-[3,4-dimethyl-5-isoxazolyl]benzenesulfonamide;		
	N'-[3,4-Dimethyl-5-isoxazolyl]sulfanilamide)		
	Minimum 99% Crystalline		

[127-69-5] C₁₁H₁₃N₃O₃S FW 267.3

S 9526 (O-S-C)	SULFITE OXIDASE	50 units	19.75
	(Sulfite: oxygen oxidoreductase; EC 1.8.3.1)	250 units	75.35
	From Chicken Liver		
	Suspension in 3.2 M ammonium sulfate containing 1.6 mM molybdic acid, pH 7.5.		
	Activity: 20-70 units per mg protein (Biuret). Unit Definition: One unit will oxidize 1.0 μ mole of sulfite to sulfate during the reduction of cytochrome c per min at pH 8.5 at 25°C.		

[9029-38-3]

p-SULFOBENZENEAZO-o-SULFOBENZENEAZO- β -NAPHTHOL-3,6-DISULFONIC ACID SODIUM SALT

See: Ponceau S Page 854